CENTRO DI RADIOLOGIA
E FISIOTERAPIA S.R.L.
Accreditato A.T.S.
Via Roma, 28 - 24020 Gorle (BG)
Cod.Fisc. 01509260160
Gruppo IVA Maredo - P. IVA 04786280166
Cap. Soc. € 500.000,00 i.v. - REA di Bergamo n. 212289
Tel. 035/290636 - 035/4236140 - Fax 035/290358
www.centroradiofisio.it - info@centroradiofisio.it
Direttore Sanitario: Dott. R. Suardi

COLONSCOPIC PREPARATION WITH PLENVU FOR NON-FAVIC PATIENTS

(6-phosphate dehydrogenase deficiency)

Three days before the exam: do not eat legumes, fruits, vegetables and foods containing seeds (e.g., sesame bread) In case of constipation (hard stools less frequently than 3 times a week) add in the 5 days before the exam, two sachets per day of MOVICOL 25 ML oral solution sachets.

The day before the exam: you may eat breakfast followed by a light meal (noodle soup or low-fat yogurt without fruit) to be completed by 2 p.m. You may drink in the afternoon, but dinner is not allowed. Do not drink alcohol, milk or anything red or purple colored or any other beverage containing pulp.
Breakfast is not allowed on the day of the exam, however, clear liquid consumption is allowed up to 3 hours before the examination.

NB Since the product tastes very sweet, at the patient's discretion, ACE (Orange, carrot and lemon) fruit juice (without pulp) may be added to make the solution more palatable (up to 150-200 cc per 500 cc PLENVU solution).

For the success of the examination, it is imperative to strictly carry out the indications written below.

For any warnings in the use of the preparation or contraindications, read the package leaflet or consult your physician. Please come to the clinic 30' before your appointment time. Come accompanied: patients are strongly advised against driving vehicles for at least 8 hours after narcosis. Follow the preparation.

- Colonoscopy scheduled between 8 a.m. and 2 p.m.

The day before the examination at 8 - 8.30 p.m. drink dose 1 of PLENVU within 30 minutes. Then drink AT LEAST another ½ litre of clear liquids (still water, tea, camomile). Take a break of one hour and drink dose 2 of PLENVU within 30 minutes. Then drink AT LEAST another ½ litre of clear liquids (still water, tea, camomile tea). In the morning, drinking water is permitted up to two hours before the start of the examination.

- Colonoscopy scheduled after 2 p.m.

The day before the examination at 8-8.30 p.m. drink dose 1 of PLENVU within 30 minutes. Then drink AT LEAST another ½ litre of clear liquids (still water, tea, camomile). On the day of the test: 4 hours before the start of the test, drink dose 2 of PLENVU within 30 minutes. Then drink AT LEAST another ½ litre of clear liquids (still water, tea, camomile). Two hours before the start of the examination, stop drinking all liquids.

How to prepare the two doses of **PLENVU**:

Dose 1

At least 1 hour must pass between the end of dose 1 and the start of dose 2.

Dose 2